

ROFENAID 40 - sulfadimethoxine and ormetoprim powder

Alpharma Inc. Animal Health

Sulfadimethoxine & Ormetoprim

Type A Medicated Article

(medicated premix)

Antibacterial

Anticoccidial

For use in broiler, replacement (breeder and layer) chickens, turkeys, ducks and chukar partridges only.

ACTIVE DRUG INGREDIENTS

Sulfadimethoxine 113.5 grams per pound (25%), and ormetoprim 68.1 grams per pound (15%) in a carrier suitable for incorporation in feed.

For use in manufacturing medicated feeds

BROILER AND REPLACEMENT (BREEDER AND LAYER) CHICKENS

Indications

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*; as an aid in the prevention of bacterial infections caused by *Haemophilus gallinarum* (infectious coryza), *Escherichia coli* (colibacillosis) and *Pasteurella multocida* (fowl cholera).

Use directions

Add 1 lb of RofenAid 40 to one ton of feed to provide 0.02% (0.0125% sulfadimethoxine and 0.0075% ormetoprim) concentration in feed.

Limitations

Feed continuously as the sole ration.

Warning

Do not feed to chickens over 16 weeks (112 days) of age.

TURKEYS

Indications

As an aid in the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. gallopavonis*, and *E. meleagrimitis*; for the prevention of bacterial infections caused by *Pasteurella multocida* (fowl cholera).

Use directions

Add 1/2 lb of RofenAid 40 to one ton of feed to provide 0.01% (0.00625% sulfadimethoxine and 0.00375% ormetoprim) concentration in feed.

Limitations

Feed continuously as the sole ration.

DUCKS

Indications

As an aid in the control of *Pasteurella multocida* (fowl cholera) in ducks including breeding ducks.

Use directions

Add 2 lbs of RofenAid 40 to one ton of feed to provide 0.04% (0.025% sulfadimethoxine and 0.015% ormetoprim) concentration in feed.

Indications

As an aid in the control of *Escherichia coli*, *Riemerella anatipestifer* (strains 2,3 & 4) and severe challenge of *P. multocida* (fowl cholera).

Use directions

Add 4 lbs of RofenAid 40 to one ton of feed to provide 0.08% (0.050% sulfadimethoxine and 0.030% ormetoprim) concentration in feed.

Limitations

Feed for 7 days as the sole ration.

Caution

For control of duck disease, medication should be started at the first signs of infection. The safety of feeding RofenAid 40 at levels higher than 0.04% in breeding ducks has not been established.

CHUKAR PARTRIDGES**Indications**

For the prevention of coccidiosis caused by *Eimeria kofoidi* and *E. legionensis*.

Use directions

Add 1 lb of RofenAid 40 to one ton of feed to provide 0.02% (0.0125% sulfadimethoxine and 0.0075% ormetoprim) concentration in feed.

Limitations

Feed continuously as the sole ration to young birds up to 8 weeks of age.

The safety of feeding RofenAid 40 to breeding stock has not been established.

WARNING

Do not feed to turkeys and ducks producing eggs for food. Withdraw 5 days prior to slaughter of broiler and replacement chickens, turkeys and ducks.

Note: Manufacture of Type B or C feeds from this product requires a Medicated Feed License Application approved by the FDA. Labels for feeds containing RofenAid 40 must contain appropriate indications, limitations and warnings as well as required feed ingredient information.

Restricted Drug (California) - Use only as directed. Not for human use.

NADA 40-209, Approved by FDA

Marketed by

Alpharma Inc.

Bridgewater, New Jersey 08807

Net wt 50 LB (22.68 kg)

**Made in USA**

Trademarks

registered by

Alpharma Inc.

770302 0910

PRINCIPAL DISPLAY PANEL - 50 LB BAG

RofenAid®

40

ALPHARMA

Sulfadimethoxine & Ormetoprim

Type A Medicated Article

(medicated premix)

Antibacterial

Anticoccidial

Important: See back panel for warnings and use directions

Net wt 50 LB (22.68 kg)

ALPHARMA®

RofenAid®

40

